

Non-Coronary Arterial Disease (VII)

Internal Carotid Artery Revascularization

Jean-Louis Mas

Neurology Department and Stroke Unit, Hôpital Sainte-Anne, Paris, France

Carotid endarterectomy remains the standard revascularization technique for the prevention of ischemic stroke resulting from severe carotid stenosis. Surgery is highly beneficial in patients with a symptomatic stenosis of 70% or greater that is not a total or near-total occlusion. The benefit becomes more diluted in patients with a symptomatic 50%-69% stenosis, and surgery has no effect, or even increases the risk of stroke, in those with a less than 50% stenosis. Surgery has also been shown to reduce the risk of stroke in asymptomatic patients with a 60%-99% stenosis, but the absolute benefit is only 1% per year. There is no clear evidence that surgery benefits asymptomatic women at 5-year follow-up, and the benefit is unknown in asymptomatic patients aged over 75 years. Decision-making must take individual factors into consideration, particularly in patients with an asymptomatic (60%-99%) or a moderate (50%-69%) symptomatic carotid stenosis, so that the risk-benefit ratio of surgery can be optimized.

Current data do not support the preferential use of carotid stenting over carotid endarterectomy in patients with a symptomatic or asymptomatic carotid stenosis who are good candidates for surgery. In those who are not good surgical candidates, carotid stenting might be equivalent to surgery, but whether or not any form of carotid revascularization is superior to medical treatment alone remains unknown.

Key words: Carotid arteries. Atherosclerosis. Clinical trials. Revascularization. Stent. Surgery. Cerebral ischemia.

Revascularización de la arteria carótida interna

La endarterectomía de la arteria carótida continúa siendo el tratamiento de revascularización estándar para la prevención de los accidentes cardiovasculares isquémicos derivados de una estenosis carotídea severa. En los pacientes con una estenosis sintomática del 70% o superior, pero sin oclusión total o subtotal de la arteria, la cirugía ofrece grandes ventajas. Éstas son más dudosas en los pacientes con estenosis sintomática del 50-69%, y la cirugía no tiene efecto alguno, e incluso aumenta el riesgo de accidente cerebrovascular, en aquellas personas cuya estenosis es inferior al 50%. La intervención quirúrgica también ha demostrado reducir el riesgo de accidentes cerebrovasculares en pacientes asintomáticos con una estenosis del 60-99%, pero el beneficio absoluto es sólo del 1% por año. No hay pruebas evidentes del beneficio de la cirugía en mujeres asintomáticas tras 5 años de seguimiento y se desconocen las ventajas de ésta en pacientes asintomáticos mayores de 75 años. Antes de tomar decisiones al respecto se deben considerar los factores individuales, en particular en pacientes con estenosis carotídea asintomática (60-99%) o moderadamente sintomática (50-69%), con el fin de mejorar la relación beneficio/riesgo de la cirugía.

Los datos actuales no avalan la preferencia de la angioplastia con endoprótesis sobre la endarterectomía en los pacientes con estenosis carotídea sintomática o asintomática que sean buenos candidatos para la cirugía. En malos candidatos quirúrgicos, la endoprótesis puede ser equivalente a la cirugía, pero aún queda por determinar si cualquiera de los tipos de revascularización de la carotídea es superior al tratamiento médico aislado.

Palabras clave: Arterias carótidas. Aterosclerosis. Ensayos clínicos. Revascularización. Endoprótesis. Cirugía. Isquemia cerebral.

Section Sponsored by Laboratorio Dr Esteve

Correspondence: Prof. Jean-Louis Mas,
Service de Neurologie, Hôpital Sainte-Anne,
1 rue Cabanis, 75674 Paris Cedex 14, France,
E-mail: jl.mas@ch-sainte-anne.fr

ABBREVIATIONS

CAS: carotid angioplasty and stenting
 ACAS: Asymptomatic Carotid Atherosclerosis Study
 CABG: coronary artery bypass graft
 CEA: carotid endarterectomy
 ECST: European Carotid Surgery Trial
 ICA: internal carotid artery
 RCTs: randomized clinical trials

INTRODUCTION

Atherosclerosis is a systemic disease which involves large- and medium-sized arteries and commonly affects the carotid artery at its bifurcation. The prevalence of cervical internal carotid artery (ICA) stenosis greater than 50% increases from about 0.5% in people aged 50-59, to 10% in those more than 80-years-old.¹ Twenty percent to 30% of patients with prior myocardial infarction or symptomatic peripheral vascular disease have >60% asymptomatic ICA stenosis.²

Carotid stenosis is an important risk factor of ischemic stroke. Approximately 10% of patients presenting with stroke have underlying carotid stenosis of 50% or more.^{3,4} This definition probably underestimates the contribution of carotid atherosclerosis to stroke⁵ but allows a standardized approach. The primary mechanism of stroke in patients with carotid artery stenosis is embolism of atherosclerotic debris or thrombotic material from the plaque into the distal cerebral vasculature. Infiltration of inflammatory cells to the surface of carotid plaques may be a critical step in promoting plaque rupture and resultant embolization, or carotid occlusion.⁶ Hemodynamic factors are correlated with increased stroke risk in patients with carotid stenosis.⁷

Cervical-artery atherosclerosis is a strong and independent predictor of myocardial infarction and vascular death; this risk increases with the degree of stenosis.^{8,9} In a recent population health survey, the annual death rate (mainly from cardiovascular disease) was 3.4% in patients with carotid stenosis detected by ultrasound screening compared with 1.2% in subjects without stenosis.^{8,9} The aims of prevention are to limit the progression of atherosclerosis and to decrease the risk of stroke and other vascular events. Therefore, carotid revascularization is only a part of the treatment of patients with carotid artery disease, which also includes strict management of risk factors and antiplatelet therapy. Carotid revascularization mainly involves carotid endarterectomy (CEA). As compared with CEA, carotid angioplasty and stenting (CAS) avoids the need for general anesthesia and an incision in the neck that could lead to nerve injury and wound complications. The costs may be less than those of surgery, mainly because the hospital stay is shorter. However, stenting also carries a risk of stroke and local complications, and the long-term efficacy of this technique is not well known. CAS is still being evaluated in the setting of randomized clinical trials (RCTs). The indications for carotid revascularization mainly depend on the degree of stenosis and whether the patient has recently experienced neurological symptoms related to the carotid artery.

ASYMPTOMATIC CAROTID STENOSIS

Natural History

Longitudinal studies in patient cohorts with carotid ultrasound suggested that the risk of first-ever ipsilateral stroke increases with the degree of stenosis; the risk is about 2% yearly in patients with stenosis greater than

TABLE 1. Risk of First-Ever Ipsilateral Stroke According to the Degree of Stenosis in Patients With Carotid Stenosis*

	Number of Patients	Mean Follow-Up, Years	Degree of Stenosis, %	Annual Risk of Ipsilateral Stroke, %
ECST	1270	4.5	0-29	1.8
	843		30-69	2.1
	127		70-99	5.7
	55		Occlusion	3.7
NASCET	1496	5	<50	1.6
	108		50-59	2.6
	113		60-74	3.0
	74		75-94	3.7
	29		95-99	2.9
	86		Occlusion	1.9

*From Inzitari et al¹² and The European Carotid Surgery Trialists Collaborative Group.¹³

50%.^{10,11} Analysis of the natural history risk of stroke from asymptomatic contralateral stenosis in RCTs of CEA for symptomatic carotid stenosis^{12,13} (Table 1), or from medical arms in RCTs of CEA for asymptomatic carotid stenosis,^{14,15} yielded a similar annual risk of 2% in patients with 50% stenosis or greater. The risk of ischemic stroke seems to decrease in patients with a very high degree ("near occlusion") of stenosis (Table 1).^{12,13} suggesting that low post-stenotic flow may protect the brain from infarction by reducing the frequency of cerebral embolism.¹⁶ Strokes occurring in the territory of an asymptomatic carotid stenosis are not always caused by the carotid lesion. Indeed, data from North American Symptomatic Carotid Endarterectomy Trial (NASCET) show that almost 50% of the strokes in the territory of a greater than 60% asymptomatic carotid stenosis are due to small vessel disease ("lacunae") and cardioembolic disease.¹² In addition to the degree of stenosis, a rapid progression of stenosis,¹⁷ features of unstable carotid lesions by ultrasound,¹⁸ cerebral hemodynamic impairment,¹⁹ the presence of silent brain infarcts,¹⁰ and a high frequency of micro-emboli at transcranial Doppler^{20,21} have been associated with an increased risk of stroke. Results of studies, however, are conflicting.

Although available information is scarce, patients who present with asymptomatic carotid occlusion have a low-risk of subsequent ischemic stroke. In a study including 30 asymptomatic patients with carotid occlusion followed for 34 months, only 1 experienced a stroke which was in another territory.^{5,22} Stroke in the territory of carotid occlusion seems to result mainly from compromised cerebral blood flow. Other potential mechanisms include emboli arising from the distal or proximal ICA stump, or from plaques in the common ICA, or external carotid artery. These find their way to the ipsilateral hemisphere or retina via collateral pathways involving the external carotid artery.²³

Carotid Endarterectomy

Guidelines on CEA for asymptomatic carotid stenosis are largely based on the results of 2 large, randomized studies of CEA versus best medical therapy, Asymptomatic Carotid Atherosclerosis Study (ACAS),¹⁴ and Asymptomatic Carotid Surgery Trial (ACST).¹⁵ In ACAS,¹⁴ a total of 1662 patients with asymptomatic carotid artery stenosis of 60% or greater reduction in diameter were randomized. ACAS only accepted surgeons with an excellent safety record: 40% of applicants were rejected and some surgeons were excluded during the trial because of adverse operative outcomes. The aggregate risk over 5 years for ipsilateral stroke and any perioperative stroke or death was estimated to be 5.1% for surgical patients, and 11.0% for patients treated medically (aggregate risk reduction of 53% [95% confidence interval, 22%-

72%]). In the ACST,¹⁵ 3120 patients with over 60% mainly asymptomatic carotid stenosis (12% had symptoms at least 6 months previously) were randomized to immediate CEA plus medical treatment versus medical treatment alone or until the operation became necessary. Surgeons were required to provide evidence of an operative risk of 6% or less for their last 50 patients having a CEA for asymptomatic stenosis, but none were excluded on the basis of his or her operative risk during the trial. Selection of patients was based on the uncertainty principle, with very few exclusion criteria. The net 5-year risks were 6.4% versus 11.8% for all strokes (net gain, 5.4% [3.0-7.8]; $P < .0001$) and 3.5% versus 6.1% for fatal or disabling strokes (net gain, 2.5% [0.8-4.3]; $P = .004$). Thus, despite differences in methods, the results are remarkably similar. The absolute reduction in 5-year risk of stroke (or any operative death) with surgery was about 5%, whereas the absolute reduction in 5-year risk of disabling or fatal stroke was about 2.5%. About 100 patients would need to undergo the operation to prevent 1 event per year and that about 200 patients would need to be treated to avoid 1 disabling or fatal stroke per year. The main differences between the trials were in the combined operative risk of stroke and death (1.5%; 95% CI, 0.6%-2.4%; in ACAS vs 3.0%; 95% CI, 2.1%-4.0%, in ACST). While the low operative risks in ACAS may not be matched in routine clinical practice, the operative risks in ACST were similar to those reported in a recent systematic review of surgical case series.²⁴

It should be stressed that, even in the optimal environment of clinical trials, the absolute risk reduction of stroke with CEA was only about 1% per year. The crossover of the Kaplan-Meier curves occurs between 1 and 2 years after surgery and the benefit of CEA becomes significant only after 5 years of follow-up. Therefore, CEA should not be performed in centres with surgical complication rates $>3\%$ or in patients with less than a 5-year life-expectancy.

As the overall benefit of CEA is marginal in asymptomatic patients, specific subgroups of patients with a higher benefit-risk ratio of surgery need to be identified. In a pooled analysis of the data from ACAS and ACST,²⁴ there was clear evidence for benefit from surgery in men, but not in women at 5 years follow-up, although some benefit may accrue with longer follow-up in ACST.²⁴ Benefit from CEA in asymptomatic patients over the age of 75 years is unknown. Patients over the age of 80 years were excluded from ACAS.¹⁴ No significant net benefit of successful CEA was found among the 650 patients older than 75 years of age randomized in ACST.¹⁵ This finding might well be a false negative result. However, these patients have a short life expectancy (half of them die within 5 years from unrelated causes) and any net benefits would probably be of limited duration.

In contrast to trials of CEA in patients with symptomatic carotid stenosis, neither ACST¹⁵ nor ACAS¹⁴ showed increasing benefit from surgery with increasing degree of stenosis within the 60%-99% range. Other potential approaches to identify patients at high risk of stroke and more likely to benefit from surgery include transcranial Doppler to identify patients with microembolic signals,²¹ hemodynamic assessment to identify patients with inadequate collateral supply,⁷ and imaging methods to identify unstable atherosclerotic plaques.²⁵

Carotid Stenting

There is no data to support the preferential use of CAS over CEA in asymptomatic patients that are good surgical candidates. The only randomized trial²⁶ has been limited to patients at high risk for surgical complications. Any conclusion regarding efficacy of CAS compared to CEA can be drawn from nonrandomized registries.^{27,28} Regarding safety, periprocedural stroke and death rates in these registries were over 3%, a complication rate that would obviate any benefit with CAS for asymptomatic patients, based on the rates observed in the ACAS and ACST surgical trials.

CAS has been generally limited to patients with perceived high risk for perioperative morbidity and mortality, despite very little evidence supporting its value. Factors commonly associated with higher surgical risk have included octogenarians, significant cardiac, pulmonary, or renal comorbidities, and anatomic factors such as surgically inaccessible lesions, a stenosis that recurred after CEA or was caused by radiation therapy, contralateral stenosis, or occlusion. Many of these factors were exclusion criteria for the large surgical endarterectomy trials. It must be stressed that these factors are associated with a higher risk for complications from CEA, not a higher risk for stroke on medical therapy. The natural history on medical therapy for patients with asymptomatic carotid stenosis and risk factors for a poor surgical outcome is unknown.^{27,28}

The Stenting and Angioplasty with Protection in Patients at High Risk for Endarterectomy (SAPPHIRE) study²⁶ was a randomized noninferiority trial comparing CEA to CAS in patients at high risk for surgery. There was no medical therapy control group. Most patients (71%) were asymptomatic ($\geq 80\%$ stenosis). The cumulative incidence of death, stroke, or myocardial infarction within 30 days after the intervention or death, or ipsilateral stroke between 31 days and 1 year occurred in 12.2% of patients randomly assigned to undergo CAS and in 20.1% of those randomly assigned to undergo CEA, leading to the conclusion that among patients with severe carotid-artery stenosis and coexisting conditions, carotid stenting with the use of an emboli-protection device is not inferior to carotid endarterectomy. One of the major issues with this trial was the absence of a

medical control group, precluding any conclusion on the superiority of CEA or CAS over medical treatment alone, in this particular population. If the risk of stroke on medical therapy in these patients at high risk for surgery is similar to the stroke risk in patients that are good surgical candidates, there would be no benefit with carotid revascularization (either by CAS or CEA), given the high rates of 30-day and 1-year outcomes in both the CAS, and CEA arms in SAPPHIRE. Therefore, the safety and efficacy of CAS and CEA in patients at high risk for surgery need to be tested against medical therapy.^{27,28}

SYMPTOMATIC CAROTID STENOSIS

Natural History

The risk of recurrent ipsilateral stroke in patients with carotid artery stenosis is much higher than the risk of a first-ever stroke and is clearly related to the severity of carotid stenosis.²⁹⁻³¹ In NASCET,^{29,30} the risk of recurrent ipsilateral stroke was 3.7% per year in patients with less than 50% carotid stenosis. This risk rose to 4.4% in patients with 50%-69% stenosis and to 13% in those with greater than 70% stenosis. However, this increased risk of stroke was present mainly during the 2 to 3 years after the first ischemic event. Then, the risk returns to a low baseline. In addition to the degree of stenosis, plaque surface irregularity is an independent risk factor for stroke.^{32,33} In a pooled analysis of European Carotid Surgery Trial (ECST) and NASCET,³⁴ the risk of ipsilateral ischemic stroke fell with time since last event, rose with age, and was higher in men than in women, higher in patients presenting with hemispheric events than retinal events, and in diabetics. Other risk factors include neurological signs, severity of the atheromatous disease,³⁵ and cerebral hemodynamic compromise.¹⁹

Patients with symptomatic ICA occlusion have an overall risk of subsequent stroke of approximately 5.5% per year and a risk of ipsilateral stroke of 2.1% per year.³⁶ Some studies suggest that the risk of recurrent stroke is increased in patients with impaired hemodynamic measurements.^{36,37}

Carotid Endarterectomy

Findings from 2 large randomized, clinical trials—NASCET and ECST^{30,31}—have established endarterectomy as the standard treatment for severe symptomatic carotid-artery stenosis. Patients were recruited if they had had a recent carotid distribution transient ischemic attack, non-disabling ischemic stroke, or a retinal infarction, in the territory of a stenosed carotid artery. Before randomization, patients were seen by a neurologist or stroke physician to confirm their eligibility, and that the symptomatic carotid artery was imaged by





Figure 1. Methods of measurement of the degree of stenosis on an angiogram in NASCET and ECST.

selective catheter angiography. In both studies, patients were stratified according to the severity of stenosis based on catheter angiography measurements. Figure 1 shows methods of measurement of the degree of carotid stenosis on an arterial angiogram in NASCET and ECST. In both studies, the group of patients with most benefit from CEA was the group with severe stenosis (70%-99%), in whom ipsilateral fatal and disabling strokes were all

significantly reduced by more than 60% over a 2-year period. The benefit became more diluted or disappeared with more moderate degrees of stenosis and in patients with near occlusions (Figure 2).

In a recent meta-analysis³⁸ of individual patients' data from the main randomized trials, using the same method of measurement of stenosis (NASCET) and definitions of outcome events, surgery increased the 5-year risk of

Figure 2. Effect of CEA on absolute risk of ipsilateral ischemic stroke (and any operative stroke or death) at 3, 5, and 8 years' follow-up by degree of symptomatic carotid stenosis, in analysis of pooled data from ECST and NASCET. From Rothwell et al.³⁸ ARR indicates absolute risk reduction.



ipsilateral ischemic stroke in patients with less than 30% stenosis (n=1746, absolute risk reduction -2.2%, $P=.05$), had no effect in patients with 30%-49% stenosis (n=1429, 3.2%, $P=.6$), was of marginal benefit in those with 50%-69% stenosis (n=1549, 4.6%, $P=.04$), but the benefit increased over 5 and 8 years, and was highly beneficial in those with 70% stenosis or greater without near-occlusion (n=1095, 16.0%, $P<.001$). There was a trend towards benefit from surgery in patients with near-occlusion at 2 years' follow-up (n=262, 5.6%, $P=.19$), but no benefit at 5 years (-1.7%, $P=.9$).

Some features alter the benefit-to-risk ratio for CEA for moderate carotid stenosis. Benefits were greatest among those with more severe stenosis, those ≥ 75 years of age, men, patients with recent stroke (rather than TIA), and patients with hemispheric symptoms rather than transient monocular blindness.^{30,39}

In a pooled analysis of ECST and NASCET trials,³⁴ benefit from surgery was greatest in men, patients aged 75 years or older, and those randomized within 2 weeks after their last ischemic event, and fell rapidly with increasing delay. For patients with 50% or higher stenosis, the number of patients needed to undergo surgery (ie, number needed to treat) to prevent 1 ipsilateral stroke in 5 years was 9 for men versus 36 for women, 5 for age 75 years or older versus 18 for younger than 65 years, and 5 for those randomized within 2 weeks after their last ischemic event, versus 125 for patients randomized after more than 12 weeks. These results were consistent across the 50%-69% and 70% or more stenosis groups, and across the 2 trials. Therefore, in patients with carotid TIA or nondisabling ischemic strokes, the procedure should be done within 2 weeks of the patient's last symptoms.³⁴ Other radiographic factors found to predict better outcomes after CEA include the presence of intracranial stenosis, the absence of leukoaraiosis, and the presence of collaterals.^{30,40,41} Table 2 shows the recommendations from the American Heart Association/American Stroke Association Council on Stroke for Interventional Approaches to Patients With Stroke Caused by Large-Artery Atherosclerotic Disease.⁴² In a systematic review of all studies published from 1980 to 2000 that reported the risk of stroke and death resulting from CEA, the operative risk was 5.1% (4.6-5.6) for symptomatic stenosis.⁴³

Carotid Stenting

In 2005, a systematic review⁴⁴ of 5 randomized trials comparing stenting with endarterectomy concluded that the current evidence does not support a change from the recommendation of carotid endarterectomy as the standard treatment for carotid stenosis. No significant difference in the major risks of treatment was found but the wide confidence intervals indicate that it is not possible to exclude a difference in favor of 1 treatment. The 30-day risks of stroke or death were 8.1% for patients treated with CAS and 6.3% for those treated

with CEA; the odds ratios for treatment-related death or any stroke after 30 days were 1.33 (95% CI, 0.86-2.04), and 1.01 (95% CI, 0.71-1.44) after 1 year. The authors recommended that patients suitable for carotid endarterectomy should only be offered stenting within the ongoing randomized trials of stenting versus surgery.

Since this time, the short-term results of the Stent-Protected Angioplasty versus Carotid Endarterectomy in symptomatic patients (SPACE)⁴⁵ and Endarterectomy Versus Stenting in patients with Symptomatic Severe carotid Stenosis (EVA-3S)⁴⁶ trials have been published, almost doubling the number of patients available for analysis.

EVA-3S is a French multicentre, non-inferiority randomized trial with national research organization funding to compare stenting with endarterectomy in patients with a symptomatic carotid stenosis of at least 60%.⁴⁶ Patients were eligible if they had experienced a carotid TIA or nondisabling stroke within 4 months before randomization, associated with an atherosclerotic stenosis within the ipsilateral carotid bifurcation of at least 60% NASCET, which investigators believe was suitable for both carotid surgery and angioplasty. The interventional physicians had to document at least 12 cases of CAS. Carotid angioplasty initially consisted of stenting with or without the use of cerebral protection. After 80 patients had been treated in the CAS arm, the Safety Committee recommended using cerebral protection devices systematically because the 30-day rate of stroke was 3.9 (0.9 to 16.7) times higher than that of unprotected CAS. The trial was stopped prematurely after the inclusion of 527 patients for reasons of both safety and futility. The 30-day incidence of any stroke or death was 3.9% after endarterectomy (95% CI, 2.0-7.2) and 9.6% after stenting (95% CI, 6.4-14.0); the relative risk of any stroke or death after stenting as compared with endarterectomy was 2.5 (95% CI, 1.2-5.1). The 30-day incidence of disabling stroke or death was 1.5% after endarterectomy (95% CI, 0.5-4.2) and 3.4% after stenting (95% CI, 1.7-6.7); the relative risk was 2.2 (95% CI, 0.7-7.2). At 6 months, the incidence of any stroke or death was 6.1% after endarterectomy and 11.7% after stenting ($P=.02$). In this study of patients with symptomatic carotid stenosis of 60% or more, the rates of death and stroke at 1 and 6 months were lower with endarterectomy than with stenting.

SPACE⁴⁵ also investigated whether CAS is not inferior to CEA in patients over the age of 50 years with symptomatic (transient ischemic attack or minor stroke) stenosis (at least 70% ECST) eligible for both methods. The primary investigator of each centre had to demonstrate 25 successful interventions prior to participation in the trial, whereas secondary investigators got a preliminary certificate after 10 interventions. In all, 1200 patients were randomized between March 2001 and January 2006. Of the CAS patients 26.6% were

TABLE 2. Recommendations from the American Heart Association/American Stroke Association Council on Stroke for Interventional Approaches to Patients With Stroke Caused by Extracranial Carotid Disease*

Recommendation	Class, Level, of Evidence
For patients with recent TIA or ischemic stroke within the last 6 months and ipsilateral severe (70%-99%) carotid artery stenosis, CEA is recommended by a surgeon with a perioperative morbidity and mortality of <6%	Class I, level A
For patients with recent TIA or ischemic stroke and ipsilateral moderate (50%-69%) carotid stenosis, CEA is recommended, depending on patient-specific factors such as age, gender, comorbidities, and severity of initial symptoms	Class I, level A
When degree of stenosis is <50%, there is no indication for CEA	Class III, level A
When CEA is indicated, surgery within 2 weeks rather than delayed surgery is suggested	Class IIa, level B
Among patients with symptomatic severe stenosis (>70%) in whom the stenosis is difficult to access surgically, medical conditions are present that greatly increase the risk for surgery, or when other specific circumstances exist such as radiation-induced stenosis or restenosis after CEA, CAS is not inferior to endarterectomy and may be considered	Class IIb, level B
CAS is reasonable when performed by operators with established periprocedural morbidity and mortality rates of 4%-6%, similar to that observed in trials of CEA and CAS	Class IIa, level B
Among patients with symptomatic carotid occlusion, EC/IC bypass surgery is not routinely recommended	Class III, level A

*Class I: conditions for which there is evidence for and/or general agreement that the procedure or treatment is useful and effective.

Class II: conditions for which there is conflicting evidence and/or a divergence of opinion about the usefulness/efficacy of a procedure or treatment.

Class IIa: weight of evidence or opinion is in favor of the procedure or treatment.

Class IIb: usefulness/efficacy is less well established by evidence or opinion.

Class III: conditions for which there is evidence and/or general agreement that the procedure or treatment is not useful/effective and in some cases may be harmful.

Level of evidence A: data derived from multiple randomized clinical trials.

Level of evidence B: data derived from a single randomized trial or nonrandomized studies.

Level of evidence C: expert opinion or case studies.

From Sacco et al.⁴²

treated using an embolic protection device. The rate of ipsilateral stroke and death between randomization and day 30 after treatment (primary endpoint) was 6.84% in the CAS and 6.34% in the CEA group. As the upper limit of the 90% confidence interval of the absolute risk reduction exceeded the predefined non-inferiority margin of 2.5%, SPACE failed to prove the non-inferiority of CAS compared with CEA in treating patients with symptomatic carotid artery stenosis. The authors concluded that the results of this trial do not justify the widespread use in the short-term of carotid-artery stenting for treatment of carotid-artery stenoses.

When comparable endpoints are used, both studies show similar results regarding the 30 day rate of stroke or death with carotid stenting: 7.7% (95% CI, 5.7%-10.1%) in SPACE compared to 9.6% (95% CI, 6.3%-13.8%) in EVA-3S, with a large overlap between confidence intervals. If we focus on the 30 day rate of severe events (ie, disabling stroke or death), it was apparently higher in SPACE (4.8%; 95% CI, 3.4%-

6.8%) than in EVA-3S (3.4%; 95% CI, 1.6%-6.4%), but once again there is a large overlap between confidence intervals. The 30 day rate of stroke or death in the surgical group was lower in EVA-3S (3.9%) than in SPACE (6.5%), resulting in a larger difference between the stenting and surgery groups, which partly explains why there is a statistically significant difference in favour of surgery in EVA-3S, but not in SPACE. However, the lower 30 day rate of stroke or death in the surgical arm of EVA-3S may be due to chance, since the confidence interval of this rate (3.9%; 95% CI, 1.9%-7.0%) includes the value of 6.5% reported in SPACE.⁴⁷

An updated meta-analysis⁴⁸ of all studies shows a significant 41% relative increase of any stroke and death within 30 days after treatment in the endovascular treatment group compared to the surgical group (OR, 1.41; 95% CI, 1.07-1.87; $P=.016$). However, a significant heterogeneity was found in this analysis. Possible reasons for the heterogeneity are the facts that some trials included asymptomatic patients, and

that the endovascular technique changed during the time period under consideration. There was a non significant 33% relative increase of disabling stroke and death after treatment in the endovascular treatment group compared to the surgical group (OR, 1.33; 95% CI, 0.89-1.98; $P=.17$). No significant heterogeneity was found for this analysis. According to present knowledge, CAS is not safer than CEA, nor does it provide a better outcome, at least in the short term (30 days). Long-term follow-up data of the SPACE and EVA-3S trials are not yet available. In particular, the rates of restenosis after surgical and endovascular treatment may have an impact on decision making—despite restenosis being asymptomatic in most cases. Patients should be informed that surgical treatment is still the standard treatment for prevention of recurrent ipsilateral ischemic stroke resulting from severe carotid stenosis.

A large number of patients will be needed to identify which factors related to patient characteristics, operator experience, and the procedure itself are associated with a high risk of stroke after CAS. In this respect, operator experience was not a major determinant of the 30-day risk of stroke or death in the EVA-3S trial. Among the patients in the stenting group, 15.8% were treated by interventional physicians who had performed more than 50 carotid-stenting procedures, 45.4% by physicians who had performed 50 or fewer procedures, and 38.8% by physicians still in procedural training. The 30-day risk of stroke or death for these 3 groups was 12.2%, 11.0%, and 7.1%, respectively ($P=.49$).⁴⁷

In symptomatic patients considered at high risk for surgery, data from recent registries of CAS show much higher 30-day rates of stroke or death than those reported in the SAPHIRE trial.²⁶ The recently published multicentre registry Carotid RX Acculink/Accunet Post-Approval Trial to Uncover Unanticipated or Rare Events (CAPTURE),⁴⁹ for example, reported a 30-day risk of stroke, death or myocardial infarction of 12% among 483 patients with symptomatic stenosis at high risk for surgery.

CAD and Stroke After Coronary Artery Bypass

In a systematic review of the literature from 1970 to 2000,⁵⁰ the risk of stroke after coronary artery bypass graft (CABG) was 2%, with a mortality of 23%. The risk increased to 3% among patients with predominantly asymptomatic unilateral carotid stenosis of 50%-99%, 5% among those with bilateral stenoses of 50%-99%, and 7%-11% among those with carotid occlusion. Significant predictive factors for post-CABG stroke included; *a*) carotid bruit (odds ratio 3.6; 95% CI, 2.8-4.6), *b*) prior stroke/TIA (odds ratio 3.6; 95% CI, 2.7-4.9), and *c*) severe carotid stenosis/occlusion (odds ratio 4.3; 95% CI, 3.2-5.7). This systematic review indicated that 50% of stroke sufferers did not have significant

carotid disease and 60% of territorial infarctions on CT scan/autopsy could not be attributed to carotid disease alone.

Most strokes in patients undergoing cardiac surgery, including those with carotid stenosis, are not related to hypoperfusion.^{50,51} In 1 study, only 9% of strokes after CABG were in watershed (hypoperfusion) areas.⁵¹ Most perioperative strokes in such patients cannot be attributed to carotid stenosis because of their location (contralateral to the affected carotid artery or bilateral). Most perioperative strokes result from manipulations of the heart and aorta, or release of particulate matter from the cardiopulmonary-bypass pump. Delayed embolism is often attributed to postoperative atrial fibrillation, myocardial infarction and hypercoagulability due to surgical trauma, and associated tissue injury. General anesthesia, dehydration, bed rest, stasis in the postoperative period, and perioperative withholding of antiplatelet or anticoagulant agents can aggravate surgery-induced hypercoagulability. Other, less common causes of perioperative stroke include air, fat, or paradoxical embolism, and arterial dissection resulting from neck manipulations during induction of anesthesia and neck surgery.⁵²

The benefit of prophylactic carotid surgery or stenting to prevent peri-operative strokes in patients undergoing CABG or other major vascular surgery remains unproven. Revascularization before surgery is generally unwarranted, but some patients with hemodynamically significant, high-grade, asymptomatic carotid stenosis—in particular those with bilateral stenoses—may benefit from carotid revascularization before elective surgery.⁵⁰⁻⁵³ Another debate is whether CEA should be staged or synchronous. In a systematic review of 97 published studies following 8972 staged or synchronous operations, 10%-12% of patients undergoing staged or synchronous procedures suffered death or major cardiovascular morbidity (stroke, MI) within 30 days of surgery. Overall, there was no significant difference in outcomes for staged and synchronous procedures, and no comparable data for patients with combined cardiac and carotid disease not undergoing staged or synchronous surgery.⁵⁴ Randomized clinical trials are needed.

CONCLUSION AND FUTURE DIRECTIONS

Although CEA is highly beneficial in patients with symptomatic severe (70% or more) carotid stenosis, it is of marginal benefit in patients with asymptomatic or symptomatic but moderate carotid disease. Consequently, specific subgroups of patients with a higher benefit – risk ratio of surgery need to be identified. Characterization of plaques in terms of morphology (eg, fibrous cap rupture) and inflammatory constituents by modern imaging techniques (eg, high-resolution MRI), identification of clinically silent emboli by transcranial Doppler and of inadequate collateral supply by hemodynamic assessment,

and biological markers may help identify unstable plaques and patients at high risk of first or recurrent stroke and allow the development of effective preventive and therapeutic measures.

Current data do not support the preferential use of CAS over CEA in patients that are good surgical candidates. In those that are not good surgical candidates, any revascularization procedure must be proven superior to medical therapy alone. Several RCTs are in progress or are being organized to compare CAS with CEA in patients at low or high risk for surgery and with symptomatic or asymptomatic carotid stenosis. CAS must be limited to

these trials designed to provide this proof of efficacy and participation to these trials must be encouraged. Nonrandomized registry data will not provide any useful information to help guide therapeutic decisions.

Among several other important areas for further investigation pertaining to CEA, the benefits and risks of CEA in octogenarians, the management of coexisting carotid and coronary artery disease, and whether improved medical treatment for atherosclerosis (eg, statins) may erase the small benefit of CE in patients with asymptomatic stenosis or 50% to 69% symptomatic stenosis need to be further explored.

REFERENCES

1. Warlow CP. Surgical treatment of asymptomatic carotid stenosis. *Cerebrovasc Dis.* 2006;6:7-14.
2. Derdeyn CP, Powers WJ. Cost-effectiveness of screening for asymptomatic carotid atherosclerotic disease. *Stroke.* 1996;27:1944-50.
3. Grau AJ, Weimar C, Bugge F, Heinrich A, Goertler M, Neumaier S, et al. Risk Factors, Outcome, and Treatment in Subtypes of Ischemic Stroke: The German Stroke Data Bank. *Stroke.* 2001;32:2559-66.
4. Bogousslavsky J, Van Melle G, Regli F. The Lausanne Stroke Registry: analysis of 1,000 consecutive patients with first stroke. *Stroke.* 1988;19:1083-92.
5. Wasserman BA, Wityk RJ, Trout HH, III, Virmani R. Low-grade carotid stenosis: looking beyond the lumen with MRI. *Stroke.* 2005;36:2504-13.
6. Golledge J, Greenhalgh RM, Davies AH. The symptomatic carotid plaque. *Stroke.* 2000;31:774-81.
7. Silvestrini M, Vernieri F, Pasqualetti P, Matteis M, Passarelli F. Impaired cerebral vasoreactivity and risk of stroke in patients with asymptomatic carotid artery stenosis. *JAMA.* 2000;283:2122-7.
8. Norris JW, Zhu CZ, Bornstein NM, Chambers BR. Vascular risks of asymptomatic carotid stenosis. *Stroke.* 1991;22:1485-90.
9. Joakimsen O, Bonna KH, Mathiesen EB, Stensland-Bugge E, Arnesen E. Prediction of mortality by ultrasound screening of a general population for carotid stenosis: the Tromso Study. *Stroke.* 2000;31:1871-6.
10. Norris JW, Halliday A. Is ultrasound sufficient for vascular imaging prior to carotid endarterectomy? *Stroke.* 2004;35:370-1.
11. Longstreth WT Jr, Bernick C, Fitzpatrick A, et al. Frequency and predictors of stroke death in 5,888 participants in the Cardiovascular Health Study. *Neurology.* 2001;56:368-75.
12. Inzitari D, Eliasziw M, Gates P, Sharpe BL, Chan RKT, Meldrum HE, et al. The causes and risk of stroke in patients with asymptomatic internal-carotid-artery stenosis. North American Symptomatic Carotid Endarterectomy Trial Collaborators. *N Engl J Med.* 2000;342:1693-700.
13. Risk of stroke in the distribution of an asymptomatic carotid artery. The European Carotid Surgery Trialists Collaborative Group. *Lancet.* 1995;345:209-12.
14. Endarterectomy for asymptomatic carotid artery stenosis. Executive Committee for the Asymptomatic Carotid Atherosclerosis Study. *JAMA.* 1995;273:1421-8.
15. MRC Asymptomatic Carotid Surgery Trial (ACST) Collaborative Group. Prevention of disabling and fatal strokes by successful carotid endarterectomy in patients without recent neurological symptoms: randomised controlled trial. *Lancet.* 2004;363:1491-502.
16. Rothwell PM, Warlow CP. Low risk of ischemic stroke in patients with reduced internal carotid artery lumen diameter distal to severe symptomatic carotid stenosis: cerebral protection due to low poststenotic flow? On behalf of the European Carotid Surgery Trialists' Collaborative Group. *Stroke.* 2000;31:622-30.
17. Hennerici M, Hulsbomer HB, Hefter H, Lammerts D, Rautenberg W. Natural history of asymptomatic extracranial arterial disease. Results of a long-term prospective study. *Brain.* 1987;110:777-91.
18. Arnold JA, Modaresi KB, Thomas N, Taylor PR, Padayachee TS. Carotid plaque characterization by duplex scanning: observer error may undermine current clinical trials. *Stroke.* 1999;30:61-5.
19. Derdeyn CP, Grubb RL Jr, Powers WJ. Cerebral hemodynamic impairment: methods of measurement and association with stroke risk. *Neurology.* 1999;53:251-9.
20. Molloy J, Markus HS. Asymptomatic embolization predicts stroke and TIA risk in patients with carotid artery stenosis. *Stroke.* 1999;30:1440-3.
21. Spence JD, Tamayo A, Lownie SP, Ng WP, Ferguson GG. Absence of microemboli on transcranial Doppler identifies low-risk patients with asymptomatic carotid stenosis. *Stroke.* 2005;36:2373-8.
22. Powers WJ, Derdeyn CP, Fritsch SM et al. Benign prognosis of never-symptomatic carotid occlusion. *Neurology.* 2000;54:878-82.
23. Klijn CJ, Kappelle LJ, Tulleken CA, Van GJ. Symptomatic carotid artery occlusion. A reappraisal of hemodynamic factors. *Stroke.* 1997;28:2084-93.
24. Rothwell PM, Goldstein LB. Carotid endarterectomy for asymptomatic carotid stenosis: asymptomatic carotid surgery trial. *Stroke.* 2004;35:2425-7.
25. Touze E, Toussaint JF, Coste J, et al. Reproducibility of high-resolution MRI for the identification and the quantification of carotid atherosclerotic plaque components: consequences for prognosis studies and therapeutic trials. *Stroke.* 2007;38:1812-9.
26. Yadav JS, Wholey MH, Kuntz RE, Schmitt E. Protected carotid-artery stenting versus endarterectomy in high-risk patients. *N Engl J Med.* 2004;351:1493-501.
27. Fayad P. Endarterectomy and stenting for asymptomatic carotid stenosis: a race at breakneck speed. *Stroke.* 2007;38 2 Suppl:707-14.
28. Derdeyn CP. Carotid stenting for asymptomatic carotid stenosis: trial it. *Stroke.* 2007;38 2 Suppl:715-20.
29. North American Symptomatic Carotid Endarterectomy Trial. Methods, patient characteristics, and progress. *Stroke.* 1991;22:711-20.
30. Barnett HJ, Taylor DW, Eliasziw M, Fox AJ, Ferguson GG, Haynes RB, et al. Benefit of carotid endarterectomy in patients with symptomatic moderate or severe stenosis. North American Symptomatic Carotid Endarterectomy Trial Collaborators. *N Engl J Med.* 1998;339:1415-25.
31. Randomised trial of endarterectomy for recently symptomatic carotid stenosis: final results of the MRC European Carotid Surgery Trial (ECST). *Lancet.* 1998;351:1379-87.
32. Rothwell PM, Gibson R, Warlow CP. Interrelation between plaque surface morphology and degree of stenosis on carotid angiograms and the risk of ischemic stroke in patients with symptomatic carotid stenosis. On behalf of the European Carotid Surgery Trialists' Collaborative Group. *Stroke.* 2000;31:615-21.
33. Eliasziw M, Streifler JY, Fox AJ, Hachinski VC, Ferguson GG, Barnett HJ. Significance of plaque ulceration in symptomatic patients with high-grade carotid stenosis. North American Symptomatic Carotid Endarterectomy Trial. *Stroke.* 1994;25:304-8.
34. Rothwell PM, Eliasziw M, Gutnikov SA, Warlow CP, Barnett HJ. Endarterectomy for symptomatic carotid stenosis in relation to clinical subgroups and timing of surgery. *Lancet.* 2004;363:915-24.
35. Rothwell PM, Warlow CP. Prediction of benefit from carotid endarterectomy in individual patients: a risk-modelling study. European Carotid Surgery Trialists' Collaborative Group. *Lancet.* 1999;353:2105-10.
36. Klijn CJ, Kappelle LJ, Tulleken CA, Van Gijn J. Symptomatic carotid artery occlusion. A reappraisal of hemodynamic factors. *Stroke.* 1997;28:2084-93.
37. Vernieri F, Pasqualetti P, Passarelli F, Rossini PM, Silvestrini M. Outcome of carotid artery occlusion is predicted by cerebrovascular reactivity. *Stroke.* 1999;30:593-8.
38. Rothwell PM, Eliasziw M, Gutnikov SA, Fox AJ, Taylor DW, Mayberg MR, et al. Analysis of pooled data from the randomised controlled trials of endarterectomy for symptomatic carotid stenosis. *Lancet.* 2003;361:107-16.
39. Streifler JY, Eliasziw M, Benavente OR, Harbison JW, Hachinski VC, Barnett JH, et al. The risk of stroke in patients with first-ever retinal vs hemispheric transient ischemic attacks and high-grade carotid stenosis. North American Symptomatic Carotid Endarterectomy Trial. *Arch Neurol.* 1995;52:246-9.
40. Kappelle LJ, Eliasziw M, Fox AJ, Sharpe BL, Barnett HJ. Importance of intracranial atherosclerotic disease in patients with symptomatic stenosis of the internal carotid artery. The North American Symptomatic Carotid Endarterectomy Trial. *Stroke.* 1999;30:282-6.
41. Henderson RD, Eliasziw M, Fox AJ, Rothwell PM, Barnett HJ. Angiographically defined collateral circulation and risk of stroke in patients with severe carotid artery stenosis. North American Symptomatic Carotid Endarterectomy Trial (NASCET) Group. *Stroke.* 2000;31:1128.

42. Sacco RL, Adams R, Albers G, Albers MJ, Benavente O, Furie K, et al. Guidelines for prevention of stroke in patients with ischemic stroke or transient ischemic attack: a statement for healthcare professionals from the American Heart Association/American Stroke Association Council on Stroke: Co-Sponsored by the Council on Cardiovascular Radiology and Intervention: the American Academy of Neurology affirms the value of this guideline. *Stroke*. 2006;37:577-617.
43. Bond R, Rerkasem K, Rothwell PM. Systematic review of the risks of carotid endarterectomy in relation to the clinical indication for and timing of surgery. *Stroke*. 2003;34:2290-301.
44. Coward LJ, Featherstone RL, Brown MM. Safety and efficacy of endovascular treatment of carotid artery stenosis compared with carotid endarterectomy: a Cochrane systematic review of the randomized evidence. *Stroke*. 2005;36:905-11.
45. Ringleb PA, Allenberg J, Bruckmann H, Branchereau A, Moulin T, Becquemin JP. 30 day results from the SPACE trial of stent-protected angioplasty versus carotid endarterectomy in symptomatic patients: a randomised non-inferiority trial. *Lancet*. 2006;68:1239-47.
46. Mas JL, Chatellier G, Beyssen B, Branchereau A, Moulin T, Becquemin JP, et al. Endarterectomy versus stenting in patients with symptomatic severe carotid stenosis. *N Engl J Med*. 2006;355:1660-71.
47. Mas JL, Chatellier G. Recent carotid stenting trials. *Lancet Neurol*. 2007;6:295-6.
48. Kern R, Ringleb PA, Hacke W, Mas JL, Hennerici MG. Stenting for carotid artery stenosis. *Nat Clin Pract Neurol*. 2007;3:212-20.
49. Gray WA, Yadav JS, Verta P, et al. The CAPTURE registry: results of carotid stenting with embolic protection in the post approval setting. *Catheter Cardiovasc Interv*. 2007;69:341-8.
50. Naylor AR, Mehta Z, Rothwell PM, Bell PR. Carotid artery disease and stroke during coronary artery bypass: a critical review of the literature. *Eur J Vasc Endovasc Surg*. 2002;23:283-94.
51. Likosky DS, Marrin CA, Caplan LR, Baribeau YR, Morton JR, Weintraub RM, et al. Determination of etiologic mechanisms of strokes secondary to coronary artery bypass graft surgery. *Stroke*. 2003;34:2830-4.
52. Selim M. Perioperative stroke. *N Engl J Med*. 2007;356:706-13.
53. Chaturvedi S, Bruno A, Feasby T, Holloway R, Benavente O, Cohen SN, et al. Carotid endarterectomy: an evidence-based review: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. 2005;65:794-801.
54. Naylor AR, Bell PR. Does the risk of post-CABG stroke merit staged or synchronous reconstruction in patients with asymptomatic carotid disease? *J Cardiovasc Surg (Torino)*. 2003;44:383-394.