

Original article

rs11613352 Polymorphism (TT Genotype) Associates with a Decrease of Triglycerides and an Increase of HDL in Familial Hypercholesterolemia Patients

Rosa Aledo,^a Teresa Padró,^a Pedro Mata,^b Rodrigo Alonso,^b and Lina Badimon^{a,c,*}^aCentro de Investigación Cardiovascular, CSIC-ICCC, IIBSant Pau, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain^bFundación Jiménez Díaz and Fundación Hipercolesterolemia Familiar, Madrid, Spain^cCátedra de Investigación Cardiovascular-Universidad Autónoma de Barcelona, Barcelona, Spain

Article history:

Received 9 December 2013

Accepted 2 April 2014

Available online 30 September 2014

Keywords:

Single nucleotide polymorphism
High-density lipoprotein cholesterol
Triglyceride
Familial hypercholesterolemia

ABSTRACT

Introduction and objectives: Recent genome-wide association studies have identified a locus on chromosome 12q13.3 associated with plasma levels of triglyceride and high-density lipoprotein cholesterol, with rs11613352 being the lead single nucleotide polymorphism in this genome-wide association study locus. The aim of the study is to investigate the involvement of rs11613352 in a population with high cardiovascular risk due to familial hypercholesterolemia.

Methods: The single nucleotide polymorphism was genotyped by Taqman[®] assay in a cohort of 601 unrelated familial hypercholesterolemia patients and its association with plasma triglyceride and high-density lipoprotein cholesterol levels was analyzed by multivariate methods based on linear regression.

Results: Minimal allele frequency was 0.17 and genotype frequencies were 0.69, 0.27, and 0.04 for CC, CT, and TT genotypes, respectively. The polymorphism is associated in a recessive manner (TT genotype) with a decrease in triglyceride levels ($P = .002$) and with an increase in high-density lipoprotein cholesterol levels ($P = .021$) after adjusting by age and sex.

Conclusions: The polymorphism rs11613352 may contribute to modulate the cardiovascular risk by modifying plasma lipid levels in familial hypercholesterolemia patients.

© 2014 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

El polimorfismo rs11613352 (genotipo TT) se asocia con disminución de triglicéridos y aumento de HDL en pacientes con hipercolesterolemia familiar

RESUMEN

Introducción y objetivos: Estudios recientes de asociación de genoma completo han identificado un locus en el cromosoma 12q.13.3 asociado con las concentraciones plasmáticas de triglicéridos y colesterol unido a lipoproteínas de alta densidad; rs11613352 es el polimorfismo de un solo nucleótido líder en dicho locus. El objetivo del estudio es investigar la implicación de rs11613352 en una población con elevado riesgo cardiovascular por hipercolesterolemia familiar.

Métodos: Se genotipificó mediante análisis Taqman[®] el polimorfismo de un solo nucleótido en una cohorte de 601 pacientes con hipercolesterolemia familiar no relacionados, y se analizó su asociación genética con las concentraciones plasmáticas de triglicéridos y colesterol unido a lipoproteínas de alta densidad, mediante métodos multivariados basados en regresión lineal.

Resultados: La frecuencia alélica mínima fue de 0,17 y las frecuencias genotípicas, 0,69, 0,27 y 0,04 para los genotipos CC, CT y TT respectivamente. El polimorfismo se asoció de manera recesiva (genotipo TT) con disminución de los triglicéridos ($p = 0,002$) y aumento de colesterol unido a lipoproteínas de alta densidad ($p = 0,021$) tras ajustar por edad y sexo.

Conclusiones: El polimorfismo rs11613352 puede contribuir a modular el riesgo cardiovascular al modificar las concentraciones plasmáticas de lípidos en los pacientes con hipercolesterolemia familiar.

© 2014 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Polimorfismo de un solo nucleótido
Colesterol unido a lipoproteínas de alta densidad
Triglicéridos
Hipercolesterolemia familiar

* Corresponding author: Centro de Investigación Cardiovascular, CSIC-ICCC, IIBSant Pau, Hospital de la Santa Creu i Sant Pau, Sant Antoni Maria Claret 167, 08025 Barcelona, Spain.

E-mail address: lbadimon@csic-iccc.org (L. Badimon).

Abbreviations

CVD: cardiovascular disease
 FH: familial hypercholesterolemia
 GWAS: genome-wide association studies
 HDL: high-density lipoproteins
 LDL: low-density lipoproteins
 TG: triglycerides

INTRODUCTION

Familial hypercholesterolemia (FH) is among the most common inherited disorders and is characterized by severely elevated plasma low-density lipoprotein (LDL) cholesterol levels and premature atherosclerotic disease. Mutations of the classical LDL receptor gene are the main cause of this disease. However, there is a substantial variation in the onset and severity of atherosclerosis in heterozygous FH patients, which might be due to environmental, metabolic, and additional genetic factors.^{1,2}

Among candidate genes/regions involved in lipid metabolism, the single nucleotide polymorphism rs11613352 found in genome-wide association studies (GWAS) as a lead single nucleotide polymorphism in the locus on chromosome 12q13.3 was associated with a significant decrease in triglycerides (TG) and with an increase in high-density lipoproteins (HDL) plasma levels³ as well as with lower serum urate levels.⁴

The aim of the study was to analyze for the first time the association of this polymorphism with plasma lipid levels in this atherosclerosis-prone population (FH).

METHODS

Population Studied

The 601 unrelated patients were randomly selected index cases of FH families from the Spanish FH Longitudinal Cohort Study (SAFEHEART) for inclusion in the study. The main characteristics of the FH Cohort have been previously reported.^{5,6} Anthropometric data such as body mass index (Kg/m²), waist circumference, waist-to-hip ratio calculated by dividing waist circumference by hip circumference, and presence of arcus cornealis, xanthomas, diabetes mellitus, hypertension, and smoking habit, along with biochemical levels of plasma lipids were obtained for each subject as previously described^{5,6} and are more deeply detailed in the [supplementary material](#).

The FH patients were considered positive for clinical manifestations of cardiovascular disease (CVD) if they had a documented history of myocardial infarction, coronary artery bypass graft surgery, percutaneous transluminal coronary angioplasty, angina pectoris with angiographically coronary atherosclerosis (>50% stenosis), ischemic atherothrombotic stroke, or chronic arterial peripheral vascular disease.

All patients included in the study were heterozygous carriers for mutations in the LDL receptor (*LDLR* gene).⁷ The biochemical and clinical characteristics of the FH population are shown in [Table 1](#).

Genotyping

DNA was obtained from whole blood by using the Kit Wizard DNA Purification from Promega. The polymorphism rs11613352 was genotyped using an allele-specific Taqman[®] assay on the Applied Biosystems[®] 7900 Real-Time PCR System. The allelic

Table 1
 Baseline Characteristics of Studied Familial Hypercholesterolemia Population

	All (n=601)	Women (n=288)	Men (n=313)	P
Age, years	49.0 (47.8-50.1)	49.7 (52.8-55.5)	48.1 (42.6-45.4)	.236
BMI, kg/m ²	27.1 (26.7-27.5)	26.0 (25.8-27.2)	27.3 (27.1-28.1)	.006
WC, cm	89.8 (87.5-90)	82.3 (80.5-84)	94.3 (92.8-95.9)	<.001
WTH	0.87 (0.86-0.88)	0.81 (0.81-0.83)	0.92 (0.91-0.94)	<.001
CVD, %	113 (18.8)	32 (11.4)	81 (25.8)	<.001
Current smokers, %	121 (20.1)	51 (17.7)	69 (22.04)	.153
Hypertension, %	99 (16.4)	62 (21.6)	37 (11.7)	.001
Diabetes mellitus, %	23 (3.8)	13 (4.5)	10 (3.2)	.524
Arcus cornealis, %	241(40.1)	106 (36.8)	135 (43.1)	.137
Xanthomas, %	126 (21.0)	60 (20.9)	66 (21.2)	.99
TC, mg/dL	245.8 (240.6-251.3)	256.6 (248.8-264.4)	235.9 (228.8-243.2)	<.001
LDL, mg/dL	177.6 (172.5-182.7)	184.2 (176.6-191.7)	171.8 (164.9-178.7)	.011
HDL, mg/dL	48.9 (47.8-49.9)	54.1 (52.8-55.5)	44 (42.6-45.4)	<.001
TG, mg/dL*	89.2 (86.1-92.3)	82.3 (78.3-86.1)	95.8 (91.6-100.3)	<.001
TC/HDL	5.0 (5.2-5.6)	4.7 (4.7-5.2)	5.4 (5.5-6.1)	<.001
LDL/HDL	3.9 (3.8-4.1)	3.6 (3.4-3.8)	4.4 (4.0-4.6)	<.001
ApoA-I, mg/dL	138.9 (136.6-141.1)	147.5 (144.5-150.9)	130.9 (127.9-133.9)	<.001
ApoB, mg/dL	117.5 (114.5-120.4)	117.8 (113.7-121.9)	117.2 (113.03-121.4)	.755

ApoA, apolipoprotein A; ApoB, apolipoprotein B; BMI, body mass index; CVD, cardiovascular disease; HDL, high-density lipoprotein; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglycerides; WC, waist circumference; WTH, waist-to-hip ratio.

Data are expressed as mean (95% confidence interval) or No. (%).

Statistical significance was tested by Student t test or Fisher exact test, respectively.

Statistical significance was set at P < .05.

* Geometric mean (95% confidence interval), with statistical significance tested by Student t on log-transformed data.

discrimination of the alleles was done using an ABI PRISM® 7900 as for genotyping.

Statistical Analysis

Quantitative data were expressed as the mean (standard deviation) and categorical data by percentages. Triglyceride values were logarithmically transformed for statistical analysis. Statistical significance was assessed by Student *t* test for quantitative data and by Fisher exact test or chi-square for categorical data.

Allele frequencies were calculated from the genotypes of the subjects. The genotype distribution in the population was assessed for Hardy-Weinberg equilibrium by chi-square test. Quantitative differences in lipid levels between genotypes were adjusted for age and sex, tested by linear regression, and summarized by means, standard errors, mean differences respect to a reference category, and 95% confidence interval (95%CI) of the differences. Various models of inheritance (general model or codominant, dominant, and recessive) were tested and genetic effects were calculated for each genotype compared to the reference genotype in each model. Statistical analysis was done using SPSS 19.0 (SPSS Inc.; Chicago, Illinois, United States) and SNPStats.⁸ *P* values < .05 were considered significant. The statistical power to detect significant differences among the genotype groups in plasma lipid levels were analyzed by Power Calculator (Quantor v2.1).

RESULTS

Clinical and Biochemical Characteristics of Studied Familial Hypercholesterolemia Population

The baseline characteristics of the FH (48% women, 52% men) population studied is shown in Table 1. The majority of patients (87%) were on lipid-lowering therapy at study inclusion and there were no differences in treatment across genotype groups (data not shown).

Association of rs11613352 With Triglycerides and High-density Lipoproteins Levels

The observed genotypic frequencies were consistent with Hardy-Weinberg equilibrium, (*P* = .32). The frequency of C and T alleles was 0.83 and 0.17, respectively in FH. The frequency of CC, CT and TT genotypes was 69%, 27% and 4%, respectively.

There were no differences in genotypes frequencies by sex or antropometric traits such as waist circumference, waist-to-hip ratio, or body mass index, nor with history of CVD, current smoking habit, or lipid levels other than TG and HDL (Student *t* test, *P* = .003 and *P* = .007, respectively). Interestingly, however, no genotype TT was observed in patients with presence of xanthomas (*P* = .012, Fisher exact test) (Table 1 of the supplementary material).

Genotype association with plasma lipid levels was only significant with TG and HDL as previously described³ and is shown in Tables 2 and 3, respectively. The best association is reached with a recessive inheritance model for plasma TG and HDL levels. Crude analysis of genetic association showed a significant association of TT genotype with a decrease in TG levels (*P* = .003), and an increase in plasma HDL levels (*P* = .0074). The association remains significant after adjusting by age and sex, *P* = .0026, mean difference of -0.29 (95%CI = -0.48 to -0.10) for natural logarithm TG values, corresponding to a change of 22 mg/dL between genotypes (Table 2) and *P* = .021, mean difference of 6.29 mg/dl (95%CI, 0.98–11.61 mg/dL) for HDL (Table 3).

Statistical power calculations assumed independence of individuals, minimal allele frequency or 0.17, a recessive genetic effect, a population mean of 4.49 and standard deviation of 0.48 for the natural logarithm of TG, mean HDL value of 48.9 mg/dL and standard deviation of 13.2 mg/dL, and a type I error of 0.05 (1-sided). With 601 FH subjects, we had 80% power to detect an effect size of -0.3 for TG, which corresponds to an approximate change of 22 mg/dL from the mean of the population, and over 70% of power to detect an effect size > 7 mg/dL from the mean HDL value.

DISCUSSION

Investigation of new GWAS candidates and their polymorphisms could help us to understand the phenotypic variability for developing clinical CVD manifestation in FH patients. Additionally, they may be of interest for future therapeutic use against atherosclerosis.

Elevated LDL, elevated TG, and decreased HDL are important risk factors for coronary artery disease^{9–11}. Not only the quantity of lipoprotein levels, but also the size and quality of lipoprotein particles are involved in atherogenesis. The particle sizes of LDL and HDL are related to plasma levels of TG and the TG/HDL ratio correlates well with the HDL particle size and with coronary artery disease.⁷ In FH, patients with elevated TG (> 150 mg/dL) display more atherogenic HDL particles with

Table 2
rs11613352 Association With Triglycerides Levels

rs11613352		Triglycerides (LN Tg) [*]	Crude analysis		Adjusted by age, sex	
Model and genotypes	No.	Response, mean (SE)	Difference (95%CI)	<i>P</i> value	Difference (95%CI)	<i>P</i> value
<i>Codominant</i>						
CC	419	4.51 (0.02)	0.00	0.011	0.00	.011
CT	161	4.49 (0.03)	-0.02 (-0.10-0.06)		-0.01 (-0.08-0.07)	
TT	21	4.22 (0.114)	-0.29 (-0.49 to -0.10)		-0.29 (-0.48 to -0.10)	
<i>Dominant</i>						
CC	419	4.51 (0.02)	0.00	0.19	0.00	.35
CT+TT	182	4.45 (0.03)	-0.05 (-0.13-0.02)		-0.04 (0.11-0.04)	
<i>Recessive</i>						
CC+CT	580	4.50 (0.02)	0.00	0.003	0.00	.0026
TT	21	4.22 (0.11)	-0.29 (-0.48 to -0.10)		-0.29 (-0.47 to -0.10)	

95%CI, 95% confidence interval; SE, standard error.

^{*} Triglyceride levels were neperian log-transformed.

Table 3
rs11613352 Association With High-density Lipoprotein Levels

rs11613352		HDL (mg/dL)	Crude analysis		Adjusted by age, sex	
Model and genotypes	No.	Response mean (SE)	Difference (95%CI)	P value	Difference (95%CI)	P value
<i>Codominant</i>						
CC	419	48.08 (0.63)	0.00	.007	0.00	.035
CT	161	50.10 (1.06)	1.02 (–0.37–4.40)		1.32 (–0.90–3.53)	
TT	21	56.50 (3.50)	8.41 (2.66–14.1)		6.66 (1.31–12.01)	
<i>Dominant</i>						
CC	419	48.08 (0.63)	0.00	.019	0.00	.076
CT+TT	182	50.83 (1.03)	2.76 (0.47–5.05)		1.93 (–0.20–4.06)	
<i>Recessive</i>						
CC+CT	580	48.64 (0.54)	0.00	.0074	0.00	.021
TT	21	56.50 (3.50)	7.85 (2.13–13.57)		6.29 (0.98–11.61)	

95%CI, 95% confidence interval; HDL, high-density lipoproteins; SE, standard error.

reduced anti-inflammatory capacity and a decreased ability to promote cholesterol efflux from macrophages, in comparison with control subjects.¹² Thus, genetic variants affecting HDL and TG may be highly relevant for the atherosclerotic process. However, the role of mutations (essentially affecting HDL levels) on CVD has been recently questioned by genetic randomization studies.¹³

By GWAS, common and rare genetic variants have been found in candidate genes contributing to dyslipidemia, although a large portion of variability remains unknown.^{14,15} The recently discovered GWAS polymorphism, rs11613352, was mainly associated with TG levels³ and also with HDL levels. The present study performed in an FH population shows for the first time that the polymorphism rs11613352 associates with variable TG and HDL levels. The TT genotype associates with the beneficial profile of lower TG and higher HDL.

The potential protective effect of the TT genotype is also emphasized by the total absence of tendon xanthomas in FH patients with TT genotype, and the presence of CVD was about half as prevalent in patients with TT genotype (Table 1 of the supplementary material), although the size of the sample did not result in statistical significance.

Several genes lie in the ±500 Kb region of the TG-suggestive GWAS signal of rs11613352.³ Among the relevant candidate genes in this region (Figure 1 of the supplementary material) are the LDL receptor-related protein 1 (*LRP1*),³ a member of the LDL-receptor family^{3,6,16}; the genes in the intergenic region *R3HDM2-INHBC-INHBE* (12q1.31–33),⁴ encoding the poorly characterized protein R3HDM2 (*KIAA1002*), an R3H domain which has been shown to bind single-stranded nucleic acid¹⁷; and *INHBC* and *INHBE* genes from the *TGF-beta* superfamily, which include among their multiple functions their roles in regulation of metabolism, homeostasis, and immune responses; and recently *ARHGAP9* have been identified as a candidate gene in adipose tissue.¹⁸ As of yet, no functional data exists linking any of the genes in proximity of rs11613352 to lipid levels.

LRP1 is a receptor involved in lipid uptake^{17,19–21} and it is localized at about 200 kb of rs11613352. In a previous study of genetic association with *LRP1* variants in FH population,⁶ we did not observe an association of 10 *LRP1* single nucleotide polymorphisms with fasting plasma lipid levels (total cholesterol, LDL, HDL, and TG) in FH (data not shown); neither have we found LD between these *LRP1* single nucleotide polymorphisms and the lead TG GWAS single nucleotide polymorphism, rs11613352; thus, we do not think that *LRP1* would be the candidate gene involved in modulating TG and HDL plasma levels.

CONCLUSIONS

We found that the polymorphism rs11613352 may contribute to attenuate cardiovascular risk by modifying plasma lipid levels in FH patients, as the TT genotype displays a profile of lower TG and higher HDL. Effects are consistent with previous large-scale GWAS results for lipid levels,^{3,4} although confirmation is needed through replication in other independent FH populations.

Rs11613352 is located within intron 1 of the poorly known gene *R3HDM2* (or *KIAA1002*) and near the transcriptionally opposite cluster *INHBC-INHBE*, a region rich in transcriptional regulatory sites. Additional expression studies are needed to elucidate the gene(s) affected by this genetic variant and its molecular function in modulating plasma lipid levels and risk of CVD.

ACKNOWLEDGEMENTS

We thank the Spanish Familial Hypercholesterolemia Foundation for assistance in the recruitment of participants and controls, and to the FH patients for their valuable contribution and willingness to participate. We would like to acknowledge Montse Gómez-Pardo for her technical assistance and the *Fundacion de Investigación Cardiovascular-Fundación Jesus Serra* for their support for the study.

FUNDING

This study was supported in part by funding from SAF (Servicio de Actividad Física) –SAF2010-16549–, CNIC (Centro Nacional de Investigaciones Cardiovasculares) –CNIC-08-2008–, RIC (Red de Investigación Cardiovascular) –RD12/0042/0027–y TerCel (Red de Terapia Celular) –RD12/0019/0026–from ISCIII (Instituto de Salud Carlos III), Spain.

CONFLICTS OF INTEREST

None declared.

SUPPLEMENTARY MATERIAL



Supplementary material associated with this article can be found in the online version available at [doi:10.1016/j.rec.2014.04.015](https://doi.org/10.1016/j.rec.2014.04.015).

REFERENCES

1. Goldstein JL, Hobbs HH, Brown MS. Familial hypercholesterolemia. In: Scriver CR, Beaudet AL, Sly WS, Valle D, editors. The metabolic and molecular bases of inherited disease. Nueva York: McGraw-Hill; 2001. p. 2863–913.
2. Jansen AC, Van Wissen S, Defesche J, Kastelein JJ. Phenotypic variability in familial hypercholesterolemia: an update. *Curr Opin Lipidol.* 2002;13:165–71.
3. Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, Kosek M, et al. Biological clinical and population relevance of 95 loci for blood lipids. *Nature.* 2010;466:707–13.
4. Yang Q, Köttgen A, Dehghan A, Smith AV, Glazer NL, Chen MH, et al. Multiple genetic loci influence serum urate levels and their relationship with gout and cardiovascular disease risk factors. *Circ Cardiovasc Genet.* 2010;3:523–30.
5. Mata N, Alonso R, Badimón L, Padró T, Fuentes F, Muñiz O, et al. Clinical Characteristics and evaluation of LDL-cholesterol treatment of the Spanish Familial Hypercholesterolemia Longitudinal Cohort Study (SAFEHEART). *Lipids Health Dis.* 2011;10:94.
6. Aledo R, Alonso R, Mata P, Llorente-Cortés V, Padró T, Badimón L. Los polimorfismos del gen *LRP1* se asocian al riesgo prematuro de enfermedad cardiovascular en pacientes con hipercolesterolemia familiar. *Rev Esp Cardiol.* 2012;65:807–12.
7. Alonso R, Defesche J, Tejedor D, Castillo S, Stef M, Mata N, et al. Genetic diagnosis of familial hypercholesterolemia using a DNA-array based platform. *Clin Biochem.* 2009;42:899–903.
8. Solé X, Guino E, Valls J, Iniesta R, Morenó V. SNPStats: a web tool for the analysis of association studies. *Bioinformatics.* 2006;22:1928–9.
9. Soska V, Jarkovsky J, Ravcukova B, Tichy L, Fajkusova L, Freiberger T. The logarithm of the triglyceride/HDL-cholesterol ratio is related to the history of cardiovascular disease in patients with familial hypercholesterolemia. *Clin Biochem.* 2012;45:96–100.
10. Baigent C, Keech A, Kearny PM, Blacwell L, Buck G, Polliciono C, et al.; Cholesterol Treatment Trialists' (CTT) Collaborators. Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins. *Lancet.* 2005;366:1267–78.
11. Sarwar N, Sandhu MS, Ricketts SL, Butterworth AS, Di Angelantonio E, Boekholdt SM, et al.; Triglyceride Coronary Disease Genetics Consortium and Emerging Risk Factors Collaboration. Triglyceride-mediated pathways and coronary disease: collaboration analysis of 101 studies. *Lancet.* 2010;375:1634–9.
12. Ottestad I, Halvorsen B, Balstad T, Otterdal K, Borge G, Brosstad F, et al. Triglyceride-rich HDL3 from patients with familial hypercholesterolemia are less able to inhibit cytokine release or to promote cholesterol efflux. *J Nutr.* 2006;136:877–81.
13. Holmes MV, Asselbergs FW, Palmer TM, Drenos F, Lanktree MB, Nelson CP, et al. Mendelian randomization of blood lipids for coronar, heart disease. *Eur Heart J.* 2014. doi: 10.1093/eurheartj/ehf571.
14. Kathiresan S, Willer CJ, Peloso GM, Demissie S, Musunuru K, Schadt EE, et al. Common variants at 30 loci contribute to polygenic dyslipidemia. *Nat Genet.* 2009;41:56–65.
15. Johansen CT, Wang J, Laktree MB, McIntyre AD, Ban MR, Martins RA, et al. An increased burden of common and rare lipid-associated risk alleles contributes to the phenotypic spectrum of hypertriglyceridemia. *Arterioscler Thromb Vasc Biol.* 2011;31:1916–26.
16. Llorente-Cortés V, Martínez-González J, Badimón L. LDL receptor related protein mediates uptake of aggregated LDL in human vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol.* 2000;20:1572–9.
17. Grishin NV. The R3H motif: a domain that binds single-stranded nucleic acids. *Trends Biochem Sci.* 1998;23:329–30.
18. Haas BE, Horvath S, Pietiläinen KH, Cantor RM, Nikkola E, Weissglas-Volkov D, et al. Adipose co-expression networks across Finns and Mexicans identify novel triglyceride-associated genes. *BMC Medical Genomics.* 2012;5:61–70.
19. Kowal RC, Hwerz J, Goldstein JL, Esser V, Brown MS. Low density lipoprotein receptor-related protein mediates uptake of cholesteryl esters derived from apoprotein E-enriched lipoproteins. *Proc Natl Acad Sci U S A.* 1989;86:5810–4.
20. Daniel TO, Shcneider WJ, Goldstein JL, Brown MS. Visualization of lipoprotein receptors by ligand blotting. *J Biol Chem.* 1983;258:4606–11.
21. Terrand J, Bruban V, Zhou L, Gong W, El Asmar Z, May P, et al. LRP1 controls intracellular cholesterol storage and fatty acid synthesis through modulation of Wnt signaling. *J Biol Chem.* 2009;284:381–8.